

Scientific Computing: An Introductory Survey

Chapter 2 – Systems of Linear Equations

Prof. Michael T. Heath

Department of Computer Science
University of Illinois at Urbana-Champaign

Copyright © 2002. Reproduction permitted
for noncommercial, educational use only.



Outline

- 1 Existence, Uniqueness, and Conditioning
- 2 Solving Linear Systems
- 3 Special Types of Linear Systems
- 4 Software for Linear Systems



Systems of Linear Equations

- Given $m \times n$ matrix A and m -vector b , find unknown n -vector x satisfying $Ax = b$
- System of equations asks “Can b be expressed as linear combination of columns of A ?”
- If so, coefficients of linear combination are given by components of solution vector x
- Solution may or may not exist, and may or may not be unique
- For now, we consider only *square case*, $m = n$



Singularity and Nonsingularity

$n \times n$ matrix \mathbf{A} is *nonsingular* if it has any of following equivalent properties

- 1 Inverse of \mathbf{A} , denoted by \mathbf{A}^{-1} , exists
- 2 $\det(\mathbf{A}) \neq 0$
- 3 $\text{rank}(\mathbf{A}) = n$
- 4 For any vector $\mathbf{z} \neq \mathbf{0}$, $\mathbf{Az} \neq \mathbf{0}$



Existence and Uniqueness

- Existence and uniqueness of solution to $Ax = b$ depend on whether A is singular or nonsingular
- Can also depend on b , but only in singular case
- If $b \in \text{span}(A)$, system is *consistent*

A	b	# solutions
nonsingular	arbitrary	one (unique)
singular	$b \in \text{span}(A)$	infinitely many
singular	$b \notin \text{span}(A)$	none



Geometric Interpretation

- In two dimensions, each equation determines straight line in plane
- Solution is intersection point of two lines
- If two straight lines are not parallel (nonsingular), then intersection point is unique
- If two straight lines are parallel (singular), then lines either do not intersect (no solution) or else coincide (any point along line is solution)
- In higher dimensions, each equation determines hyperplane; if matrix is nonsingular, intersection of hyperplanes is unique solution



Example: Nonsingularity

- 2×2 system

$$2x_1 + 3x_2 = b_1$$

$$5x_1 + 4x_2 = b_2$$

or in matrix-vector notation

$$\mathbf{Ax} = \begin{bmatrix} 2 & 3 \\ 5 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = \mathbf{b}$$

is nonsingular regardless of value of \mathbf{b}

- For example, if $\mathbf{b} = [8 \ 13]^T$, then $\mathbf{x} = [1 \ 2]^T$ is unique solution



Example: Singularity

- 2×2 system

$$\mathbf{A}\mathbf{x} = \begin{bmatrix} 2 & 3 \\ 4 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = \mathbf{b}$$

is singular regardless of value of \mathbf{b}

- With $\mathbf{b} = [4 \ 7]^T$, there is no solution
- With $\mathbf{b} = [4 \ 8]^T$, $\mathbf{x} = [\gamma \ (4 - 2\gamma)/3]^T$ is solution for any real number γ , so there are infinitely many solutions



Vector Norms

- Magnitude, modulus, or absolute value for scalars generalizes to *norm* for vectors
- We will use only p -norms, defined by

$$\|\mathbf{x}\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}$$


for integer $p > 0$ and n -vector \mathbf{x}

- Important special cases
 - 1-norm: $\|\mathbf{x}\|_1 = \sum_{i=1}^n |x_i|$
 - 2-norm: $\|\mathbf{x}\|_2 = \left(\sum_{i=1}^n |x_i|^2 \right)^{1/2}$
 - ∞ -norm: $\|\mathbf{x}\|_\infty = \max_i |x_i|$



Example: Vector Norms

- Drawing shows unit sphere in two dimensions for each norm



- Norms have following values for vector shown

$$\|\mathbf{x}\|_1 = 2.8 \quad \|\mathbf{x}\|_2 = 2.0 \quad \|\mathbf{x}\|_\infty = 1.6$$

< interactive example >



Equivalence of Norms

- In general, for any vector \mathbf{x} in \mathbb{R}^n , $\|\mathbf{x}\|_1 \geq \|\mathbf{x}\|_2 \geq \|\mathbf{x}\|_\infty$
- However, we also have

$$\|\mathbf{x}\|_1 \leq \sqrt{n} \|\mathbf{x}\|_2, \quad \|\mathbf{x}\|_2 \leq \sqrt{n} \|\mathbf{x}\|_\infty, \quad \|\mathbf{x}\|_1 \leq n \|\mathbf{x}\|_\infty$$

- Thus, for given n , norms differ by at most a constant, and hence are equivalent: if one is small, they must all be proportionally small.



Properties of Vector Norms

- For any vector norm
 - $\|x\| > 0$ if $x \neq 0$
 - $\|\gamma x\| = |\gamma| \cdot \|x\|$ for any scalar γ
 - $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality)
- In more general treatment, these properties taken as *definition* of vector norm
- Useful variation on triangle inequality
 - $|\|x\| - \|y\|| \leq \|x - y\|$



Matrix Norms

- *Matrix norm* corresponding to given vector norm is defined by

$$\|A\| = \max_{x \neq 0} \frac{\|Ax\|}{\|x\|}$$

- Norm of matrix measures maximum stretching matrix does to any vector in given vector norm



Matrix Norms

- Matrix norm corresponding to vector 1-norm is maximum absolute *column* sum

$$\|\mathbf{A}\|_1 = \max_j \sum_{i=1}^n |a_{ij}|$$

- Matrix norm corresponding to vector ∞ -norm is maximum absolute *row* sum

$$\|\mathbf{A}\|_\infty = \max_i \sum_{j=1}^n |a_{ij}|$$

- Handy way to remember these is that matrix norms agree with corresponding vector norms for $n \times 1$ matrix



Properties of Matrix Norms

- Any matrix norm satisfies
 - $\|\mathbf{A}\| > 0$ if $\mathbf{A} \neq \mathbf{0}$
 - $\|\gamma\mathbf{A}\| = |\gamma| \cdot \|\mathbf{A}\|$ for any scalar γ
 - $\|\mathbf{A} + \mathbf{B}\| \leq \|\mathbf{A}\| + \|\mathbf{B}\|$
- Matrix norms we have defined also satisfy
 - $\|\mathbf{AB}\| \leq \|\mathbf{A}\| \cdot \|\mathbf{B}\|$
 - $\|\mathbf{Ax}\| \leq \|\mathbf{A}\| \cdot \|\mathbf{x}\|$ for any vector \mathbf{x}



Condition Number

- *Condition number* of square nonsingular matrix \mathbf{A} is defined by

$$\text{cond}(\mathbf{A}) = \|\mathbf{A}\| \cdot \|\mathbf{A}^{-1}\|$$

- By convention, $\text{cond}(\mathbf{A}) = \infty$ if \mathbf{A} is singular
- Since

$$\|\mathbf{A}\| \cdot \|\mathbf{A}^{-1}\| = \left(\max_{\mathbf{x} \neq \mathbf{0}} \frac{\|\mathbf{A}\mathbf{x}\|}{\|\mathbf{x}\|} \right) \cdot \left(\min_{\mathbf{x} \neq \mathbf{0}} \frac{\|\mathbf{A}\mathbf{x}\|}{\|\mathbf{x}\|} \right)^{-1}$$

condition number measures ratio of maximum stretching to maximum shrinking matrix does to any nonzero vectors

- Large $\text{cond}(\mathbf{A})$ means \mathbf{A} is *nearly singular*



Properties of Condition Number

- For any matrix \mathbf{A} , $\text{cond}(\mathbf{A}) \geq 1$
- For identity matrix, $\text{cond}(\mathbf{I}) = 1$
- For any matrix \mathbf{A} and scalar γ , $\text{cond}(\gamma\mathbf{A}) = \text{cond}(\mathbf{A})$
- For any diagonal matrix $\mathbf{D} = \text{diag}(d_i)$, $\text{cond}(\mathbf{D}) = \frac{\max |d_i|}{\min |d_i|}$

< interactive example >



Computing Condition Number

- Definition of condition number involves matrix inverse, so it is nontrivial to compute
- Computing condition number from definition would require much more work than computing solution whose accuracy is to be assessed
- In practice, condition number is estimated inexpensively as byproduct of solution process
- Matrix norm $\|\mathbf{A}\|$ is easily computed as maximum absolute column sum (or row sum, depending on norm used)
- Estimating $\|\mathbf{A}^{-1}\|$ at low cost is more challenging



Computing Condition Number, continued

- From properties of norms, if $\mathbf{A}z = \mathbf{y}$, then

$$\frac{\|z\|}{\|y\|} \leq \|\mathbf{A}^{-1}\|$$

and bound is achieved for optimally chosen \mathbf{y}

- Efficient condition estimators heuristically pick \mathbf{y} with large ratio $\|z\|/\|y\|$, yielding good estimate for $\|\mathbf{A}^{-1}\|$
- Good software packages for linear systems provide efficient and reliable condition estimator



Error Bounds

- Condition number yields error bound for computed solution to linear system
- Let x be solution to $Ax = b$, and let \hat{x} be solution to $A\hat{x} = b + \Delta b$
- If $\Delta x = \hat{x} - x$, then

$$b + \Delta b = A(\hat{x}) = A(x + \Delta x) = Ax + A\Delta x$$

which leads to bound

$$\frac{\|\Delta x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\Delta b\|}{\|b\|}$$

for possible relative change in solution x due to relative change in right-hand side b < interactive example >



Error Bounds, continued

- Similar result holds for relative change in matrix: if $(\mathbf{A} + \mathbf{E})\hat{\mathbf{x}} = \mathbf{b}$, then

$$\frac{\|\Delta \mathbf{x}\|}{\|\hat{\mathbf{x}}\|} \leq \text{cond}(\mathbf{A}) \frac{\|\mathbf{E}\|}{\|\mathbf{A}\|}$$

- If input data are accurate to machine precision, then bound for relative error in solution \mathbf{x} becomes


$$\frac{\|\hat{\mathbf{x}} - \mathbf{x}\|}{\|\mathbf{x}\|} \leq \text{cond}(\mathbf{A}) \epsilon_{\text{mach}}$$

- Computed solution loses about $\log_{10}(\text{cond}(\mathbf{A}))$ decimal digits of accuracy relative to accuracy of input




Error Bounds – Illustration

- In two dimensions, uncertainty in intersection point of two lines depends on whether lines are nearly parallel



well-conditioned



ill-conditioned

< interactive example >



Error Bounds – Caveats

- Normwise analysis bounds relative error in *largest* components of solution; relative error in smaller components can be much larger
 - Componentwise error bounds can be obtained, but somewhat more complicated
- Conditioning of system is affected by relative scaling of rows or columns
 - Ill-conditioning can result from poor scaling as well as near singularity
 - Rescaling can help the former, but not the latter



Residual

- **Residual vector** of approximate solution \hat{x} to linear system $Ax = b$ is defined by

$$r = b - A\hat{x}$$

- In theory, if A is nonsingular, then $\|\hat{x} - x\| = 0$ if, and only if, $\|r\| = 0$, but they are not necessarily small simultaneously
- Since

$$\frac{\|\Delta x\|}{\|\hat{x}\|} \leq \text{cond}(A) \frac{\|r\|}{\|A\| \cdot \|\hat{x}\|}$$

small relative residual implies small relative error in approximate solution *only if* A is well-conditioned



Residual, continued

- If computed solution \hat{x} exactly satisfies

$$(\mathbf{A} + \mathbf{E})\hat{x} = \mathbf{b}$$

then

$$\frac{\|\mathbf{r}\|}{\|\mathbf{A}\| \|\hat{x}\|} \leq \frac{\|\mathbf{E}\|}{\|\mathbf{A}\|}$$

so large *relative residual* implies large backward error in matrix, and algorithm used to compute solution is unstable

- Stable algorithm yields small relative residual regardless of conditioning of nonsingular system
- Small residual is easy to obtain, but does not necessarily imply computed solution is accurate



Solving Linear Systems

- To solve linear system, transform it into one whose solution is same but easier to compute
- What type of transformation of linear system leaves solution unchanged?
- We can *premultiply* (from left) both sides of linear system $Ax = b$ by any *nonsingular* matrix M without affecting solution
- Solution to $MAx = Mb$ is given by

$$x = (MA)^{-1}Mb = A^{-1}M^{-1}Mb = A^{-1}b$$



Example: Permutations

- **Permutation matrix** P has one 1 in each row and column and zeros elsewhere, i.e., identity matrix with rows or columns permuted
- Note that $P^{-1} = P^T$
- Premultiplying both sides of system by permutation matrix, $PAx = Pb$, reorders rows, but solution x is unchanged
- Postmultiplying A by permutation matrix, $APx = b$, reorders columns, which permutes components of original solution

$$x = (AP)^{-1}b = P^{-1}A^{-1}b = P^T(A^{-1}b)$$



Example: Diagonal Scaling

- Row scaling: premultiplying both sides of system by nonsingular diagonal matrix D , $DAx = Db$, multiplies each row of matrix and right-hand side by corresponding diagonal entry of D , but solution x is unchanged
- Column scaling: postmultiplying A by D , $ADx = b$, multiplies each column of matrix by corresponding diagonal entry of D , which rescales original solution

$$x = (AD)^{-1}b = D^{-1}A^{-1}b$$



Triangular Linear Systems

- What type of linear system is easy to solve?
- If one equation in system involves only one component of solution (i.e., only one entry in that row of matrix is nonzero), then that component can be computed by division
- If another equation in system involves only one additional solution component, then by substituting one known component into it, we can solve for other component
- If this pattern continues, with only one new solution component per equation, then all components of solution can be computed in succession.
- System with this property is called *triangular*



Triangular Matrices

- Two specific triangular forms are of particular interest
 - *lower triangular*: all entries *above* main diagonal are zero,
 $a_{ij} = 0$ for $i < j$
 - *upper triangular*: all entries *below* main diagonal are zero,
 $a_{ij} = 0$ for $i > j$
- Successive substitution process described earlier is especially easy to formulate for lower or upper triangular systems
- Any triangular matrix can be permuted into upper or lower triangular form by suitable row permutation



Forward-Substitution

- *Forward-substitution* for lower triangular system $Lx = b$

$$x_1 = b_1/\ell_{11}, \quad x_i = \left(b_i - \sum_{j=1}^{i-1} \ell_{ij}x_j \right) / \ell_{ii}, \quad i = 2, \dots, n$$

```
for  $j = 1$  to  $n$                                 { loop over columns }  
  if  $\ell_{jj} = 0$  then stop                       { stop if matrix is singular }  
   $x_j = b_j/\ell_{jj}$                                { compute solution component }  
  for  $i = j + 1$  to  $n$   
     $b_i = b_i - \ell_{ij}x_j$                          { update right-hand side }  
  end  
end
```



Back-Substitution

- *Back-substitution* for upper triangular system $Ux = b$

$$x_n = b_n/u_{nn}, \quad x_i = \left(b_i - \sum_{j=i+1}^n u_{ij}x_j \right) / u_{ii}, \quad i = n - 1, \dots, 1$$

```
for  $j = n$  to 1                                { loop backwards over columns }  
  if  $u_{jj} = 0$  then stop                       { stop if matrix is singular }  
   $x_j = b_j/u_{jj}$                                 { compute solution component }  
  for  $i = 1$  to  $j - 1$   
     $b_i = b_i - u_{ij}x_j$                           { update right-hand side }  
  end  
end
```



Example: Triangular Linear System

$$\begin{bmatrix} 2 & 4 & -2 \\ 0 & 1 & 1 \\ 0 & 0 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 8 \end{bmatrix}$$

- Using back-substitution for this upper triangular system, last equation, $4x_3 = 8$, is solved directly to obtain $x_3 = 2$
- Next, x_3 is substituted into second equation to obtain $x_2 = 2$
- Finally, both x_3 and x_2 are substituted into first equation to obtain $x_1 = -1$



Elimination

- To transform general linear system into triangular form, we need to replace selected nonzero entries of matrix by zeros
- This can be accomplished by taking linear combinations of rows
- Consider 2-vector $\mathbf{a} = \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}$
- If $a_1 \neq 0$, then

$$\begin{bmatrix} 1 & 0 \\ -a_2/a_1 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} a_1 \\ 0 \end{bmatrix}$$



Elementary Elimination Matrices

- More generally, we can annihilate *all* entries below k th position in n -vector \mathbf{a} by transformation

$$\mathbf{M}_k \mathbf{a} = \begin{bmatrix} 1 & \cdots & 0 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 & \cdots & 0 \\ 0 & \cdots & -m_{k+1} & 1 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & -m_n & 0 & \cdots & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_k \\ a_{k+1} \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} a_1 \\ \vdots \\ a_k \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

where $m_i = a_i/a_k$, $i = k + 1, \dots, n$

- Divisor a_k , called *pivot*, must be nonzero



Elementary Elimination Matrices, continued

- Matrix M_k , called *elementary elimination matrix*, adds multiple of row k to each subsequent row, with *multipliers* m_i chosen so that result is zero
- M_k is unit lower triangular and nonsingular
- $M_k = I - m_k e_k^T$, where $m_k = [0, \dots, 0, m_{k+1}, \dots, m_n]^T$ and e_k is k th column of identity matrix
- $M_k^{-1} = I + m_k e_k^T$, which means $M_k^{-1} = L_k$ is same as M_k except signs of multipliers are reversed



Elementary Elimination Matrices, continued

- If M_j , $j > k$, is another elementary elimination matrix, with vector of multipliers m_j , then

$$\begin{aligned}M_k M_j &= I - m_k e_k^T - m_j e_j^T + m_k e_k^T m_j e_j^T \\ &= I - m_k e_k^T - m_j e_j^T\end{aligned}$$

which means product is essentially “union,” and similarly for product of inverses, $L_k L_j$



Example: Elementary Elimination Matrices

• For $\mathbf{a} = \begin{bmatrix} 2 \\ 4 \\ -2 \end{bmatrix}$,

$$\mathbf{M}_1 \mathbf{a} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \\ -2 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$$

and

$$\mathbf{M}_2 \mathbf{a} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1/2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \\ -2 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 0 \end{bmatrix}$$

Example, continued

- Note that

$$\mathbf{L}_1 = \mathbf{M}_1^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}, \quad \mathbf{L}_2 = \mathbf{M}_2^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1/2 & 1 \end{bmatrix}$$

and

$$\mathbf{M}_1\mathbf{M}_2 = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & 1/2 & 1 \end{bmatrix}, \quad \mathbf{L}_1\mathbf{L}_2 = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & -1/2 & 1 \end{bmatrix}$$



Gaussian Elimination

- To reduce general linear system $Ax = b$ to upper triangular form, first choose M_1 , with a_{11} as pivot, to annihilate first column of A below first row
 - System becomes $M_1Ax = M_1b$, but solution is unchanged
- Next choose M_2 , using a_{22} as pivot, to annihilate second column of M_1A below second row
 - System becomes $M_2M_1Ax = M_2M_1b$, but solution is still unchanged
- Process continues for each successive column until all subdiagonal entries have been zeroed



Gaussian Elimination, continued

- Resulting upper triangular linear system

$$\begin{aligned}M_{n-1} \cdots M_1 A x &= M_{n-1} \cdots M_1 b \\M A x &= M b\end{aligned}$$

can be solved by back-substitution to obtain solution to original linear system $Ax = b$

- Process just described is called *Gaussian elimination*



LU Factorization

- Product $L_k L_j$ is unit lower triangular if $k < j$, so

$$L = M^{-1} = M_1^{-1} \cdots M_{n-1}^{-1} = L_1 \cdots L_{n-1}$$

is unit lower triangular

- By design, $U = MA$ is upper triangular
- So we have

$$A = LU$$

with L unit lower triangular and U upper triangular

- Thus, Gaussian elimination produces *LU factorization* of matrix into triangular factors



LU Factorization, continued

- Having obtained LU factorization, $Ax = b$ becomes $LUx = b$, and can be solved by forward-substitution in lower triangular system $Ly = b$, followed by back-substitution in upper triangular system $Ux = y$
- Note that $y = Mb$ is same as transformed right-hand side in Gaussian elimination
- Gaussian elimination and LU factorization are two ways of expressing same solution process



Example: Gaussian Elimination

- Use Gaussian elimination to solve linear system

$$\mathbf{Ax} = \begin{bmatrix} 2 & 4 & -2 \\ 4 & 9 & -3 \\ -2 & -3 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 8 \\ 10 \end{bmatrix} = \mathbf{b}$$

- To annihilate subdiagonal entries of first column of \mathbf{A} ,

$$\mathbf{M}_1\mathbf{A} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 & -2 \\ 4 & 9 & -3 \\ -2 & -3 & 7 \end{bmatrix} = \begin{bmatrix} 2 & 4 & -2 \\ 0 & 1 & 1 \\ 0 & 1 & 5 \end{bmatrix},$$

$$\mathbf{M}_1\mathbf{b} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 8 \\ 10 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 12 \end{bmatrix}$$



Example, continued

- To annihilate subdiagonal entry of second column of $M_1\mathbf{A}$,

$$M_2M_1\mathbf{A} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 & -2 \\ 0 & 1 & 1 \\ 0 & 1 & 5 \end{bmatrix} = \begin{bmatrix} 2 & 4 & -2 \\ 0 & 1 & 1 \\ 0 & 0 & 4 \end{bmatrix} = \mathbf{U},$$

$$M_2M_1\mathbf{b} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \\ 12 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 8 \end{bmatrix} = \mathbf{Mb}$$



Example, continued

- We have reduced original system to equivalent upper triangular system

$$U\mathbf{x} = \begin{bmatrix} 2 & 4 & -2 \\ 0 & 1 & 1 \\ 0 & 0 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 8 \end{bmatrix} = M\mathbf{b}$$

which can now be solved by back-substitution to obtain

$$\mathbf{x} = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix}$$



Example, continued

- To write out LU factorization explicitly,

$$\mathbf{L}_1 \mathbf{L}_2 = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} = \mathbf{L}$$

so that

$$\mathbf{A} = \begin{bmatrix} 2 & 4 & -2 \\ 4 & 9 & -3 \\ -2 & -3 & 7 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 4 & -2 \\ 0 & 1 & 1 \\ 0 & 0 & 4 \end{bmatrix} = \mathbf{LU}$$



Row Interchanges

- Gaussian elimination breaks down if leading diagonal entry of remaining unreduced matrix is zero at any stage
- Easy fix: if diagonal entry in column k is zero, then interchange row k with some subsequent row having nonzero entry in column k and then proceed as usual
- If there is no nonzero on or below diagonal in column k , then there is nothing to do at this stage, so skip to next column
- Zero on diagonal causes resulting upper triangular matrix U to be singular, but LU factorization can still be completed
- Subsequent back-substitution will fail, however, as it should for singular matrix



Partial Pivoting

- In principle, any nonzero value will do as pivot, but in practice pivot should be chosen to minimize error propagation
- To avoid amplifying previous rounding errors when multiplying remaining portion of matrix by elementary elimination matrix, multipliers should not exceed 1 in magnitude
- This can be accomplished by choosing entry of largest magnitude on or below diagonal as pivot at each stage
- Such *partial pivoting* is essential in practice for numerically stable implementation of Gaussian elimination for general linear systems [< interactive example >](#)



LU Factorization with Partial Pivoting

- With partial pivoting, each M_k is preceded by permutation P_k to interchange rows to bring entry of largest magnitude into diagonal pivot position
- Still obtain $MA = U$, with U upper triangular, but now

$$M = M_{n-1}P_{n-1} \cdots M_1P_1$$

- $L = M^{-1}$ is still triangular in general sense, but not necessarily *lower* triangular
- Alternatively, we can write

$$PA = LU$$

where $P = P_{n-1} \cdots P_1$ permutes rows of A into order determined by partial pivoting, and now L is lower triangular



Complete Pivoting

- *Complete pivoting* is more exhaustive strategy in which largest entry in entire remaining unreduced submatrix is permuted into diagonal pivot position
- Requires interchanging columns as well as rows, leading to factorization

$$PAQ = LU$$

with L unit lower triangular, U upper triangular, and P and Q permutations

- Numerical stability of complete pivoting is theoretically superior, but pivot search is more expensive than for partial pivoting
- Numerical stability of partial pivoting is more than adequate in practice, so it is almost always used in solving linear systems by Gaussian elimination



Example: Pivoting

- Need for pivoting has nothing to do with whether matrix is singular or nearly singular
- For example,

$$\mathbf{A} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

is nonsingular yet has no LU factorization unless rows are interchanged, whereas

$$\mathbf{A} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

is singular yet has LU factorization



Example: Small Pivots

- To illustrate effect of small pivots, consider

$$\mathbf{A} = \begin{bmatrix} \epsilon & 1 \\ 1 & 1 \end{bmatrix}$$

where ϵ is positive number smaller than ϵ_{mach}

- If rows are not interchanged, then pivot is ϵ and multiplier is $-1/\epsilon$, so

$$\mathbf{M} = \begin{bmatrix} 1 & 0 \\ -1/\epsilon & 1 \end{bmatrix}, \quad \mathbf{L} = \begin{bmatrix} 1 & 0 \\ 1/\epsilon & 1 \end{bmatrix},$$

$$\mathbf{U} = \begin{bmatrix} \epsilon & 1 \\ 0 & 1 - 1/\epsilon \end{bmatrix} = \begin{bmatrix} \epsilon & 1 \\ 0 & -1/\epsilon \end{bmatrix}$$

in floating-point arithmetic, but then

$$\mathbf{LU} = \begin{bmatrix} 1 & 0 \\ 1/\epsilon & 1 \end{bmatrix} \begin{bmatrix} \epsilon & 1 \\ 0 & -1/\epsilon \end{bmatrix} = \begin{bmatrix} \epsilon & 1 \\ 1 & 0 \end{bmatrix} \neq \mathbf{A}$$



Example, continued

- Using small pivot, and correspondingly large multiplier, has caused loss of information in transformed matrix
- If rows interchanged, then pivot is 1 and multiplier is $-\epsilon$, so

$$M = \begin{bmatrix} 1 & 0 \\ -\epsilon & 1 \end{bmatrix}, \quad L = \begin{bmatrix} 1 & 0 \\ \epsilon & 1 \end{bmatrix},$$

$$U = \begin{bmatrix} 1 & 1 \\ 0 & 1 - \epsilon \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

in floating-point arithmetic

- Thus,

$$LU = \begin{bmatrix} 1 & 0 \\ \epsilon & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ \epsilon & 1 \end{bmatrix}$$

which is correct after permutation



Pivoting, continued

- Although pivoting is generally required for stability of Gaussian elimination, pivoting is *not* required for some important classes of matrices
 - *Diagonally dominant*

$$\sum_{i=1, i \neq j}^n |a_{ij}| < |a_{jj}|, \quad j = 1, \dots, n$$

- *Symmetric positive definite*

$$\mathbf{A} = \mathbf{A}^T \quad \text{and} \quad \mathbf{x}^T \mathbf{A} \mathbf{x} > 0 \quad \text{for all } \mathbf{x} \neq \mathbf{0}$$



Residual

- Residual $\mathbf{r} = \mathbf{b} - \mathbf{A}\hat{\mathbf{x}}$ for solution $\hat{\mathbf{x}}$ computed using Gaussian elimination satisfies

$$\frac{\|\mathbf{r}\|}{\|\mathbf{A}\| \|\hat{\mathbf{x}}\|} \leq \frac{\|\mathbf{E}\|}{\|\mathbf{A}\|} \leq \rho n^2 \epsilon_{\text{mach}}$$

where \mathbf{E} is backward error in matrix \mathbf{A} and *growth factor* ρ is ratio of largest entry of \mathbf{U} to largest entry of \mathbf{A}

- Without pivoting, ρ can be arbitrarily large, so Gaussian elimination without pivoting is unstable
- With partial pivoting, ρ can still be as large as 2^{n-1} , but such behavior is extremely rare



Residual, continued

- There is little or no growth in practice, so

$$\frac{\|r\|}{\|A\| \|\hat{x}\|} \leq \frac{\|E\|}{\|A\|} \approx n \epsilon_{\text{mach}}$$

which means Gaussian elimination with partial pivoting yields small relative residual regardless of conditioning of system

- Thus, small relative residual does not necessarily imply computed solution is close to “true” solution unless system is well-conditioned
- Complete pivoting yields even smaller growth factor, but additional margin of stability usually is not worth extra cost



Example: Small Residual

- Use 3-digit decimal arithmetic to solve

$$\begin{bmatrix} 0.641 & 0.242 \\ 0.321 & 0.121 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0.883 \\ 0.442 \end{bmatrix}$$

- Gaussian elimination with partial pivoting yields triangular system

$$\begin{bmatrix} 0.641 & 0.242 \\ 0 & 0.000242 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0.883 \\ -0.000383 \end{bmatrix}$$

- Back-substitution then gives solution

$$\hat{\mathbf{x}} = [0.782 \quad 1.58]^T$$

- Exact residual for this solution is

$$\mathbf{r} = \mathbf{b} - \mathbf{A}\hat{\mathbf{x}} = \begin{bmatrix} -0.000622 \\ -0.000202 \end{bmatrix}$$



Example, continued

- Residual is as small as we can expect using 3-digit arithmetic, but exact solution is

$$\mathbf{x} = [1.00 \quad 1.00]^T$$

so error is almost as large as solution

- Cause of this phenomenon is that matrix is nearly singular ($\text{cond}(\mathbf{A}) > 4000$)
- Division that determines x_2 is between two quantities that are both on order of rounding error, and hence result is essentially arbitrary
- When arbitrary value for x_2 is substituted into first equation, value for x_1 is computed so that first equation is satisfied, yielding small residual, but poor solution



Implementation of Gaussian Elimination

- Gaussian elimination has general form of triple-nested loop

```
for _____  
  for _____  
    for _____  
       $a_{ij} = a_{ij} - (a_{ik}/a_{kk})a_{kj}$   
    end  
  end  
end
```

- Indices i , j , and k of **for** loops can be taken in any order, for total of $3! = 6$ different arrangements
- These variations have different memory access patterns, which may cause their performance to vary widely on different computers



Uniqueness of LU Factorization

- Despite variations in computing it, LU factorization is unique up to diagonal scaling of factors
- Provided row pivot sequence is same, if we have two LU factorizations $PA = LU = \hat{L}\hat{U}$, then $\hat{L}^{-1}L = \hat{U}U^{-1} = D$ is both lower and upper triangular, hence diagonal
- If both L and \hat{L} are unit lower triangular, then D must be identity matrix, so $L = \hat{L}$ and $U = \hat{U}$
- Uniqueness is made explicit in LDU factorization $PA = LDU$, with L unit lower triangular, U unit upper triangular, and D diagonal



Storage Management

- Elementary elimination matrices M_k , their inverses L_k , and permutation matrices P_k used in formal description of LU factorization process are *not* formed explicitly in actual implementation
- U overwrites upper triangle of A , multipliers in L overwrite strict lower triangle of A , and unit diagonal of L need not be stored
- Row interchanges usually are not done explicitly; auxiliary integer vector keeps track of row order in original locations



Complexity of Solving Linear Systems

- LU factorization requires about $n^3/3$ floating-point multiplications and similar number of additions
- Forward- and back-substitution for single right-hand-side vector together require about n^2 multiplications and similar number of additions
- Can also solve linear system by matrix inversion:
$$\mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$$
- Computing \mathbf{A}^{-1} is tantamount to solving n linear systems, requiring LU factorization of \mathbf{A} followed by n forward- and back-substitutions, one for each column of identity matrix
- Operation count for inversion is about n^3 , three times as expensive as LU factorization



Inversion vs. Factorization

- Even with many right-hand sides \mathbf{b} , inversion never overcomes higher initial cost, since each matrix-vector multiplication $\mathbf{A}^{-1}\mathbf{b}$ requires n^2 operations, similar to cost of forward- and back-substitution
- Inversion gives less accurate answer; for example, solving $3x = 18$ by division gives $x = 18/3 = 6$, but inversion gives $x = 3^{-1} \times 18 = 0.333 \times 18 = 5.99$ using 3-digit arithmetic
- Matrix inverses often occur as convenient notation in formulas, but explicit inverse is rarely required to implement such formulas
- For example, product $\mathbf{A}^{-1}\mathbf{B}$ should be computed by LU factorization of \mathbf{A} , followed by forward- and back-substitutions using each column of \mathbf{B}



Gauss-Jordan Elimination

- In Gauss-Jordan elimination, matrix is reduced to diagonal rather than triangular form
- Row combinations are used to annihilate entries above as well as below diagonal
- Elimination matrix used for given column vector a is of form

$$\begin{bmatrix} 1 & \cdots & 0 & -m_1 & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 1 & -m_{k-1} & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 1 & 0 & \cdots & 0 \\ 0 & \cdots & 0 & -m_{k+1} & 1 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & -m_n & 0 & \cdots & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ \vdots \\ a_{k-1} \\ a_k \\ a_{k+1} \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ a_k \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

where $m_i = a_i/a_k$, $i = 1, \dots, n$



Gauss-Jordan Elimination, continued

- Gauss-Jordan elimination requires about $n^3/2$ multiplications and similar number of additions, 50% more expensive than LU factorization
- During elimination phase, same row operations are also applied to right-hand-side vector (or vectors) of system of linear equations
- Once matrix is in diagonal form, components of solution are computed by dividing each entry of transformed right-hand side by corresponding diagonal entry of matrix
- Latter requires only n divisions, but this is not enough cheaper to offset more costly elimination phase

< interactive example >



Solving Modified Problems

- If right-hand side of linear system changes but matrix does not, then LU factorization need not be repeated to solve new system
- Only forward- and back-substitution need be repeated for new right-hand side
- This is substantial savings in work, since additional triangular solutions cost only $\mathcal{O}(n^2)$ work, in contrast to $\mathcal{O}(n^3)$ cost of factorization



Sherman-Morrison Formula

- Sometimes refactorization can be avoided even when matrix *does* change
- *Sherman-Morrison formula* gives inverse of matrix resulting from rank-one change to matrix whose inverse is already known

$$(\mathbf{A} - \mathbf{u}\mathbf{v}^T)^{-1} = \mathbf{A}^{-1} + \mathbf{A}^{-1}\mathbf{u}(1 - \mathbf{v}^T\mathbf{A}^{-1}\mathbf{u})^{-1}\mathbf{v}^T\mathbf{A}^{-1}$$

where \mathbf{u} and \mathbf{v} are n -vectors

- Evaluation of formula requires $\mathcal{O}(n^2)$ work (for matrix-vector multiplications) rather than $\mathcal{O}(n^3)$ work required for inversion



Rank-One Updating of Solution

- To solve linear system $(\mathbf{A} - \mathbf{u}\mathbf{v}^T)\mathbf{x} = \mathbf{b}$ with new matrix, use Sherman-Morrison formula to obtain

$$\begin{aligned}\mathbf{x} &= (\mathbf{A} - \mathbf{u}\mathbf{v}^T)^{-1}\mathbf{b} \\ &= \mathbf{A}^{-1}\mathbf{b} + \mathbf{A}^{-1}\mathbf{u}(1 - \mathbf{v}^T\mathbf{A}^{-1}\mathbf{u})^{-1}\mathbf{v}^T\mathbf{A}^{-1}\mathbf{b}\end{aligned}$$

which can be implemented by following steps

- Solve $\mathbf{A}\mathbf{z} = \mathbf{u}$ for \mathbf{z} , so $\mathbf{z} = \mathbf{A}^{-1}\mathbf{u}$
 - Solve $\mathbf{A}\mathbf{y} = \mathbf{b}$ for \mathbf{y} , so $\mathbf{y} = \mathbf{A}^{-1}\mathbf{b}$
 - Compute $\mathbf{x} = \mathbf{y} + ((\mathbf{v}^T\mathbf{y})/(1 - \mathbf{v}^T\mathbf{z}))\mathbf{z}$
- If \mathbf{A} is already factored, procedure requires only triangular solutions and inner products, so only $\mathcal{O}(n^2)$ work and no explicit inverses



Example: Rank-One Updating of Solution

- Consider rank-one modification

$$\begin{bmatrix} 2 & 4 & -2 \\ 4 & 9 & -3 \\ -2 & -1 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 8 \\ 10 \end{bmatrix}$$

(with 3, 2 entry changed) of system whose LU factorization was computed in earlier example

- One way to choose update vectors is

$$\mathbf{u} = \begin{bmatrix} 0 \\ 0 \\ -2 \end{bmatrix} \quad \text{and} \quad \mathbf{v} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

so matrix of modified system is $\mathbf{A} - \mathbf{u}\mathbf{v}^T$



Example, continued

- Using LU factorization of A to solve $Az = u$ and $Ay = b$,

$$z = \begin{bmatrix} -3/2 \\ 1/2 \\ -1/2 \end{bmatrix} \quad \text{and} \quad y = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix}$$

- Final step computes updated solution

$$x = y + \frac{v^T y}{1 - v^T z} z = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix} + \frac{2}{1 - 1/2} \begin{bmatrix} -3/2 \\ 1/2 \\ -1/2 \end{bmatrix} = \begin{bmatrix} -7 \\ 4 \\ 0 \end{bmatrix}$$

- We have thus computed solution to modified system without factoring modified matrix



Scaling Linear Systems

- In principle, solution to linear system is unaffected by diagonal scaling of matrix and right-hand-side vector
- In practice, scaling affects both conditioning of matrix and selection of pivots in Gaussian elimination, which in turn affect numerical accuracy in finite-precision arithmetic
- It is usually best if all entries (or uncertainties in entries) of matrix have about same size
- Sometimes it may be obvious how to accomplish this by choice of measurement units for variables, but there is no foolproof method for doing so in general
- Scaling can introduce rounding errors if not done carefully



Example: Scaling

- Linear system

$$\begin{bmatrix} 1 & 0 \\ 0 & \epsilon \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 1 \\ \epsilon \end{bmatrix}$$

has condition number $1/\epsilon$, so is ill-conditioned if ϵ is small

- If second row is multiplied by $1/\epsilon$, then system becomes perfectly well-conditioned
- Apparent ill-conditioning was due purely to poor scaling
- In general, it is usually much less obvious how to correct poor scaling



Iterative Refinement

- Given approximate solution x_0 to linear system $Ax = b$, compute residual

$$r_0 = b - Ax_0$$

- Now solve linear system $Az_0 = r_0$ and take

$$x_1 = x_0 + z_0$$

as new and “better” approximate solution, since

$$\begin{aligned} Ax_1 &= A(x_0 + z_0) = Ax_0 + Az_0 \\ &= (b - r_0) + r_0 = b \end{aligned}$$

- Process can be repeated to refine solution successively until convergence, potentially producing solution accurate to full machine precision



Iterative Refinement, continued

- Iterative refinement requires double storage, since both original matrix and its LU factorization are required
- Due to cancellation, residual usually must be computed with higher precision for iterative refinement to produce meaningful improvement
- For these reasons, iterative improvement is often impractical to use routinely, but it can still be useful in some circumstances
- For example, iterative refinement can sometimes stabilize otherwise unstable algorithm



Special Types of Linear Systems

- Work and storage can often be saved in solving linear system if matrix has special properties
- Examples include
 - **Symmetric**: $A = A^T$, $a_{ij} = a_{ji}$ for all i, j
 - **Positive definite**: $x^T Ax > 0$ for all $x \neq 0$
 - **Band**: $a_{ij} = 0$ for all $|i - j| > \beta$, where β is *bandwidth* of A
 - **Sparse**: most entries of A are zero



Symmetric Positive Definite Matrices

- If A is symmetric and positive definite, then LU factorization can be arranged so that $U = L^T$, which gives *Cholesky factorization*

$$A = L L^T$$

where L is lower triangular with positive diagonal entries

- Algorithm for computing it can be derived by equating corresponding entries of A and $L L^T$
- In 2×2 case, for example,

$$\begin{bmatrix} a_{11} & a_{21} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} l_{11} & 0 \\ l_{21} & l_{22} \end{bmatrix} \begin{bmatrix} l_{11} & l_{21} \\ 0 & l_{22} \end{bmatrix}$$

implies

$$l_{11} = \sqrt{a_{11}}, \quad l_{21} = a_{21}/l_{11}, \quad l_{22} = \sqrt{a_{22} - l_{21}^2}$$



Cholesky Factorization

- One way to write resulting general algorithm, in which Cholesky factor L overwrites original matrix A , is

```
for  $j = 1$  to  $n$   
  for  $k = 1$  to  $j - 1$   
    for  $i = j$  to  $n$   
       $a_{ij} = a_{ij} - a_{ik} \cdot a_{jk}$   
    end  
  end  
   $a_{jj} = \sqrt{a_{jj}}$   
  for  $k = j + 1$  to  $n$   
     $a_{kj} = a_{kj} / a_{jj}$   
  end  
end
```



Cholesky Factorization, continued

- Features of Cholesky algorithm for symmetric positive definite matrices
 - All n square roots are of positive numbers, so algorithm is well defined
 - No pivoting is required to maintain numerical stability
 - Only lower triangle of A is accessed, and hence upper triangular portion need not be stored
 - Only $n^3/6$ multiplications and similar number of additions are required
- Thus, Cholesky factorization requires only about half work and half storage compared with LU factorization of general matrix by Gaussian elimination, and also avoids need for pivoting

< interactive example >



Symmetric Indefinite Systems

- For symmetric indefinite A , Cholesky factorization is not applicable, and some form of pivoting is generally required for numerical stability
- Factorization of form

$$PAP^T = LDL^T$$

with L unit lower triangular and D either tridiagonal or block diagonal with 1×1 and 2×2 diagonal blocks, can be computed stably using symmetric pivoting strategy

- In either case, cost is comparable to that of Cholesky factorization



Band Matrices

- Gaussian elimination for band matrices differs little from general case — only ranges of loops change
- Typically matrix is stored in array by diagonals to avoid storing zero entries
- If pivoting is required for numerical stability, bandwidth can grow (but no more than double)
- General purpose solver for arbitrary bandwidth is similar to code for Gaussian elimination for general matrices
- For fixed small bandwidth, band solver can be extremely simple, especially if pivoting is not required for stability



Tridiagonal Matrices

- Consider tridiagonal matrix

$$\mathbf{A} = \begin{bmatrix} b_1 & c_1 & 0 & \cdots & 0 \\ a_2 & b_2 & c_2 & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & a_{n-1} & b_{n-1} & c_{n-1} \\ 0 & \cdots & 0 & a_n & b_n \end{bmatrix}$$

- Gaussian elimination without pivoting reduces to

$$d_1 = b_1$$

for $i = 2$ **to** n

$$m_i = a_i / d_{i-1}$$

$$d_i = b_i - m_i c_{i-1}$$

end



Tridiagonal Matrices, continued

- LU factorization of A is then given by

$$\mathbf{L} = \begin{bmatrix} 1 & 0 & \cdots & \cdots & 0 \\ m_2 & 1 & \ddots & & \vdots \\ 0 & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & m_{n-1} & 1 & 0 \\ 0 & \cdots & 0 & m_n & 1 \end{bmatrix}, \quad \mathbf{U} = \begin{bmatrix} d_1 & c_1 & 0 & \cdots & 0 \\ 0 & d_2 & c_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ \vdots & & \ddots & d_{n-1} & c_{n-1} \\ 0 & \cdots & \cdots & 0 & d_n \end{bmatrix}$$



General Band Matrices

- In general, band system of bandwidth β requires $\mathcal{O}(\beta n)$ storage, and its factorization requires $\mathcal{O}(\beta^2 n)$ work
- Compared with full system, savings is substantial if $\beta \ll n$



Iterative Methods for Linear Systems

- Gaussian elimination is direct method for solving linear system, producing exact solution in finite number of steps (in exact arithmetic)
- Iterative methods begin with initial guess for solution and successively improve it until desired accuracy attained
- In theory, it might take infinite number of iterations to converge to exact solution, but in practice iterations are terminated when residual is as small as desired
- For some types of problems, iterative methods have significant advantages over direct methods
- We will study specific iterative methods later when we consider solution of partial differential equations



LINPACK and LAPACK

- LINPACK is software package for solving wide variety of systems of linear equations, both general dense systems and special systems, such as symmetric or banded
- Solving linear systems of such fundamental importance in scientific computing that LINPACK has become standard benchmark for comparing performance of computers
- LAPACK is more recent replacement for LINPACK featuring higher performance on modern computer architectures, including some parallel computers
- Both LINPACK and LAPACK are available from Netlib



Basic Linear Algebra Subprograms

- High-level routines in LINPACK and LAPACK are based on lower-level Basic Linear Algebra Subprograms (BLAS)
- BLAS encapsulate basic operations on vectors and matrices so they can be optimized for given computer architecture while high-level routines that call them remain portable
- Higher-level BLAS encapsulate matrix-vector and matrix-matrix operations for better utilization of memory hierarchies such as cache and virtual memory with paging
- Generic Fortran versions of BLAS are available from Netlib, and many computer vendors provide custom versions optimized for their particular systems



Examples of BLAS

Level	Work	Examples	Function
1	$\mathcal{O}(n)$	saxpy sdot snrm2	Scalar \times vector + vector Inner product Euclidean vector norm
2	$\mathcal{O}(n^2)$	sgemv strsv sger	Matrix-vector product Triangular solution Rank-one update
3	$\mathcal{O}(n^3)$	sgemm strsm ssyrk	Matrix-matrix product Multiple triang. solutions Rank- k update

- Level-3 BLAS have more opportunity for data reuse, and hence higher performance, because they perform more operations per data item than lower-level BLAS

