

## Scientific Computing: An Introductory Survey

Chapter 10 – Boundary Value Problems for  
Ordinary Differential Equations

Prof. Michael T. Heath

Department of Computer Science  
University of Illinois at Urbana-ChampaignCopyright © 2002. Reproduction permitted  
for noncommercial, educational use only.

Michael T. Heath

Scientific Computing

1 / 45

Boundary Value Problems  
Numerical Methods for BVPsBoundary Values  
Existence and Uniqueness  
Conditioning and Stability

## Boundary Value Problems

- Side conditions prescribing solution or derivative values at specified points are required to make solution of ODE unique
- For initial value problem, all side conditions are specified at single point, say  $t_0$
- For **boundary value problem** (BVP), side conditions are specified at more than one point
- $k$ th order ODE, or equivalent first-order system, requires  $k$  side conditions
- For ODEs, side conditions are typically specified at endpoints of interval  $[a, b]$ , so we have **two-point boundary value problem** with boundary conditions (BC) at  $a$  and  $b$ .



Michael T. Heath

Scientific Computing

3 / 45

Boundary Value Problems  
Numerical Methods for BVPsBoundary Values  
Existence and Uniqueness  
Conditioning and Stability

## Example: Separated Linear Boundary Conditions

- Two-point BVP for second-order scalar ODE

$$u'' = f(t, u, u'), \quad a < t < b$$

with BC

$$u(a) = \alpha, \quad u(b) = \beta$$

is equivalent to first-order system of ODEs

$$\begin{bmatrix} y_1' \\ y_2' \end{bmatrix} = \begin{bmatrix} y_2 \\ f(t, y_1, y_2) \end{bmatrix}, \quad a < t < b$$

with separated linear BC

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} y_1(a) \\ y_2(a) \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} y_1(b) \\ y_2(b) \end{bmatrix} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}$$



Michael T. Heath

Scientific Computing

5 / 45

Boundary Value Problems  
Numerical Methods for BVPsBoundary Values  
Existence and Uniqueness  
Conditioning and Stability

## Existence and Uniqueness, continued

- In general, solvability of BVP

$$\mathbf{y}' = \mathbf{f}(t, \mathbf{y}), \quad a < t < b$$

with BC

$$\mathbf{g}(\mathbf{y}(a), \mathbf{y}(b)) = \mathbf{0}$$

depends on solvability of algebraic equation

$$\mathbf{g}(\mathbf{x}, \mathbf{y}(b; \mathbf{x})) = \mathbf{0}$$

where  $\mathbf{y}(t; \mathbf{x})$  denotes solution to ODE with initial condition  $\mathbf{y}(a) = \mathbf{x}$  for  $\mathbf{x} \in \mathbb{R}^n$ 

- Solvability of latter system is difficult to establish if  $\mathbf{g}$  is nonlinear



Michael T. Heath

Scientific Computing

7 / 45

## Outline

- Boundary Value Problems
- Numerical Methods for BVPs



Michael T. Heath

Scientific Computing

2 / 45

Boundary Value Problems  
Numerical Methods for BVPsBoundary Values  
Existence and Uniqueness  
Conditioning and Stability

## Boundary Value Problems, continued

- General **first-order two-point BVP** has form

$$\mathbf{y}' = \mathbf{f}(t, \mathbf{y}), \quad a < t < b$$

with BC

$$\mathbf{g}(\mathbf{y}(a), \mathbf{y}(b)) = \mathbf{0}$$

where  $\mathbf{f}: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$  and  $\mathbf{g}: \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$ 

- Boundary conditions are **separated** if any given component of  $\mathbf{g}$  involves solution values only at  $a$  or at  $b$ , but not both
- Boundary conditions are **linear** if they are of form

$$\mathbf{B}_a \mathbf{y}(a) + \mathbf{B}_b \mathbf{y}(b) = \mathbf{c}$$

where  $\mathbf{B}_a, \mathbf{B}_b \in \mathbb{R}^{n \times n}$  and  $\mathbf{c} \in \mathbb{R}^n$ 

- BVP is **linear** if ODE and BC are both linear



Michael T. Heath

Scientific Computing

4 / 45

Boundary Value Problems  
Numerical Methods for BVPsBoundary Values  
Existence and Uniqueness  
Conditioning and Stability

## Existence and Uniqueness

- Unlike IVP, with BVP we cannot begin at initial point and continue solution step by step to nearby points
- Instead, solution is determined everywhere simultaneously, so existence and/or uniqueness may not hold
- For example,

$$u'' = -u, \quad 0 < t < b$$

with BC

$$u(0) = 0, \quad u(b) = \beta$$

with  $b$  integer multiple of  $\pi$ , has infinitely many solutions if  $\beta = 0$ , but no solution if  $\beta \neq 0$ 

Michael T. Heath

Scientific Computing

6 / 45

Boundary Value Problems  
Numerical Methods for BVPsBoundary Values  
Existence and Uniqueness  
Conditioning and Stability

## Existence and Uniqueness, continued

- For **linear** BVP, existence and uniqueness are more tractable
- Consider linear BVP

$$\mathbf{y}' = \mathbf{A}(t) \mathbf{y} + \mathbf{b}(t), \quad a < t < b$$

where  $\mathbf{A}(t)$  and  $\mathbf{b}(t)$  are continuous, with BC

$$\mathbf{B}_a \mathbf{y}(a) + \mathbf{B}_b \mathbf{y}(b) = \mathbf{c}$$

- Let  $\mathbf{Y}(t)$  denote matrix whose  $i$ th column,  $\mathbf{y}_i(t)$ , called **ith mode**, is solution to  $\mathbf{y}' = \mathbf{A}(t) \mathbf{y}$  with initial condition  $\mathbf{y}(a) = \mathbf{e}_i$ ,  $i$ th column of identity matrix

- Then BVP has unique solution if, and only if, matrix

$$\mathbf{Q} \equiv \mathbf{B}_a \mathbf{Y}(a) + \mathbf{B}_b \mathbf{Y}(b)$$

is nonsingular



Michael T. Heath

Scientific Computing

8 / 45

## Existence and Uniqueness, continued

- Assuming  $Q$  is nonsingular, define

$$\Phi(t) = Y(t) Q^{-1}$$

and **Green's function**

$$G(t, s) = \begin{cases} \Phi(t) B_a \Phi(a) \Phi^{-1}(s), & a \leq s \leq t \\ -\Phi(t) B_b \Phi(b) \Phi^{-1}(s), & t < s \leq b \end{cases}$$

- Then solution to BVP given by

$$y(t) = \Phi(t) c + \int_a^b G(t, s) b(s) ds$$

- This result also gives absolute condition number for BVP

$$\kappa = \max\{\|\Phi\|_\infty, \|G\|_\infty\}$$

I

Michael T. Heath

Scientific Computing

9 / 45

## Numerical Methods for BVPs

- For IVP, initial data supply all information necessary to begin numerical solution method at initial point and step forward from there
- For BVP, we have insufficient information to begin step-by-step numerical method, so numerical methods for solving BVPs are more complicated than those for solving IVPs
- We will consider four types of numerical methods for two-point BVPs
  - Shooting
  - Finite difference
  - Collocation
  - Galerkin

I

Michael T. Heath

Scientific Computing

11 / 45

## Shooting Method, continued

- For given  $\gamma$ , value at  $b$  of solution  $u(b)$  to IVP

$$u'' = f(t, u, u')$$

with initial conditions

$$u(a) = \alpha, \quad u'(a) = \gamma$$

can be considered as function of  $\gamma$ , say  $g(\gamma)$

- Then BVP becomes problem of solving equation  $g(\gamma) = \beta$
- One-dimensional zero finder can be used to solve this scalar equation

I

Michael T. Heath

Scientific Computing

13 / 45

## Example, continued

- We first try guess for initial slope of  $y_2(0) = 1$

$$\begin{aligned} y^{(1)} &= y^{(0)} + \frac{h}{6}(k_1 + 2k_2 + 2k_3 + k_4) \\ &= \begin{bmatrix} 0 \\ 1 \end{bmatrix} + \frac{0.5}{6} \left( \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 1.0 \\ 1.5 \end{bmatrix} + 2 \begin{bmatrix} 1.375 \\ 1.500 \end{bmatrix} + \begin{bmatrix} 1.75 \\ 3.00 \end{bmatrix} \right) = \begin{bmatrix} 0.625 \\ 1.750 \end{bmatrix} \\ y^{(2)} &= \begin{bmatrix} 0.625 \\ 1.750 \end{bmatrix} + \frac{0.5}{6} \left( \begin{bmatrix} 1.75 \\ 3.00 \end{bmatrix} + 2 \begin{bmatrix} 2.5 \\ 4.5 \end{bmatrix} + 2 \begin{bmatrix} 2.875 \\ 4.500 \end{bmatrix} + \begin{bmatrix} 4 \\ 6 \end{bmatrix} \right) = \begin{bmatrix} 2 \\ 4 \end{bmatrix} \end{aligned}$$

- So we have hit  $y_1(1) = 2$  instead of desired value  $y_1(1) = 1$

I

Michael T. Heath

Scientific Computing

15 / 45

## Conditioning and Stability

- Conditioning or stability of BVP depends on interplay between growth of solution modes and boundary conditions
- For IVP, instability is associated with modes that grow exponentially as time increases
- For BVP, solution is determined everywhere simultaneously, so there is no notion of "direction" of integration in interval  $[a, b]$
- Growth of modes increasing with time is limited by boundary conditions at  $b$ , and "growth" (in reverse) of decaying modes is limited by boundary conditions at  $a$
- For BVP to be well-conditioned, growing and decaying modes must be controlled appropriately by boundary conditions imposed

I


Michael T. Heath

Scientific Computing

10 / 45

## Shooting Method

- In statement of two-point BVP, we are given value of  $u(a)$
- If we also knew value of  $u'(a)$ , then we would have IVP that we could solve by methods discussed previously
- Lacking that information, we try sequence of increasingly accurate guesses until we find value for  $u'(a)$  such that when we solve resulting IVP, approximate solution value at  $t = b$  matches desired boundary value,  $u(b) = \beta$



I

Michael T. Heath

Scientific Computing

12 / 45

## Example: Shooting Method

- Consider two-point BVP for second-order ODE

$$u'' = 6t, \quad 0 < t < 1$$

with BC

$$u(0) = 0, \quad u(1) = 1$$

- For each guess for  $u'(0)$ , we will integrate resulting IVP using classical fourth-order Runge-Kutta method to determine how close we come to hitting desired solution value at  $t = 1$
- For simplicity of illustration, we will use step size  $h = 0.5$  to integrate IVP from  $t = 0$  to  $t = 1$  in only two steps
- First, we transform second-order ODE into system of two first-order ODEs

$$y'(t) = \begin{bmatrix} y_1'(t) \\ y_2'(t) \end{bmatrix} = \begin{bmatrix} y_2 \\ 6t \end{bmatrix}$$

I

Michael T. Heath

Scientific Computing

14 / 45

## Example, continued

- We try again, this time with initial slope  $y_2(0) = -1$

$$\begin{aligned} y^{(1)} &= \begin{bmatrix} 0 \\ -1 \end{bmatrix} + \frac{0.5}{6} \left( \begin{bmatrix} -1 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} -1.0 \\ 1.5 \end{bmatrix} + 2 \begin{bmatrix} -0.625 \\ 1.500 \end{bmatrix} + \begin{bmatrix} -0.25 \\ 3.00 \end{bmatrix} \right) \\ &= \begin{bmatrix} -0.375 \\ -0.250 \end{bmatrix} \\ y^{(2)} &= \begin{bmatrix} -0.375 \\ -0.250 \end{bmatrix} + \frac{0.5}{6} \left( \begin{bmatrix} -0.25 \\ 3.00 \end{bmatrix} + 2 \begin{bmatrix} 0.5 \\ 4.5 \end{bmatrix} + 2 \begin{bmatrix} 0.875 \\ 4.500 \end{bmatrix} + \begin{bmatrix} 2 \\ 6 \end{bmatrix} \right) \\ &= \begin{bmatrix} 0 \\ 2 \end{bmatrix} \end{aligned}$$

- So we have hit  $y_1(1) = 0$  instead of desired value  $y_1(1) = 1$ , but we now have initial slope bracketed between  $-1$  and  $1$

I

Michael T. Heath

Scientific Computing

16 / 45

## Example, continued

- We omit further iterations necessary to identify correct initial slope, which turns out to be  $y_2(0) = 0$

$$\begin{aligned} \mathbf{y}^{(1)} &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \frac{0.5}{6} \left( \begin{bmatrix} 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0.0 \\ 1.5 \end{bmatrix} + 2 \begin{bmatrix} 0.375 \\ 1.500 \end{bmatrix} + \begin{bmatrix} 0.75 \\ 3.00 \end{bmatrix} \right) \\ &= \begin{bmatrix} 0.125 \\ 0.750 \end{bmatrix} \\ \mathbf{y}^{(2)} &= \begin{bmatrix} 0.125 \\ 0.750 \end{bmatrix} + \frac{0.5}{6} \left( \begin{bmatrix} 0.75 \\ 3.00 \end{bmatrix} + 2 \begin{bmatrix} 1.5 \\ 4.5 \end{bmatrix} + 2 \begin{bmatrix} 1.875 \\ 4.500 \end{bmatrix} + \begin{bmatrix} 3 \\ 6 \end{bmatrix} \right) \\ &= \begin{bmatrix} 1 \\ 3 \end{bmatrix} \end{aligned}$$

- So we have indeed hit target solution value  $y_1(1) = 1$

Michael T. Heath

Scientific Computing

17 / 45

## Multiple Shooting

- Simple shooting method inherits stability (or instability) of associated IVP, which may be unstable even when BVP is stable
- Such ill-conditioning may make it difficult to achieve convergence of iterative method for solving nonlinear equation
- Potential remedy is **multiple shooting**, in which interval  $[a, b]$  is divided into subintervals, and shooting is carried out on each
- Requiring continuity at internal mesh points provides BC for individual subproblems
- Multiple shooting results in larger *system* of nonlinear equations to solve

Michael T. Heath

Scientific Computing

19 / 45

## Finite Difference Method, continued

- We replace derivatives by finite difference approximations such as

$$\begin{aligned} u'(t_i) &\approx \frac{y_{i+1} - y_{i-1}}{2h} \\ u''(t_i) &\approx \frac{y_{i+1} - 2y_i + y_{i-1}}{h^2} \end{aligned}$$

- This yields system of equations

$$\frac{y_{i+1} - 2y_i + y_{i-1}}{h^2} = f\left(t_i, y_i, \frac{y_{i+1} - y_{i-1}}{2h}\right)$$

to be solved for unknowns  $y_i, i = 1, \dots, n$

- System of equations may be linear or nonlinear, depending on whether  $f$  is linear or nonlinear

Michael T. Heath

Scientific Computing

21 / 45

## Example: Finite Difference Method

- Consider again two-point BVP

$$u'' = 6t, \quad 0 < t < 1$$

with BC

$$u(0) = 0, \quad u(1) = 1$$


- To keep computation to minimum, we compute approximate solution at one interior mesh point,  $t = 0.5$ , in interval  $[0, 1]$
- Including boundary points, we have three mesh points,  $t_0 = 0, t_1 = 0.5$ , and  $t_2 = 1$
- From BC, we know that  $y_0 = u(t_0) = 0$  and  $y_2 = u(t_2) = 1$ , and we seek approximate solution  $y_1 \approx u(t_1)$

Michael T. Heath

Scientific Computing

23 / 45

## Example, continued



&lt; interactive example &gt;

Michael T. Heath

Scientific Computing

18 / 45

## Finite Difference Method

- Finite difference method** converts BVP into system of algebraic equations by replacing all derivatives with finite difference approximations
- For example, to solve two-point BVP

$$u'' = f(t, u, u'), \quad a < t < b$$

with BC

$$u(a) = \alpha, \quad u(b) = \beta$$

we introduce mesh points  $t_i = a + ih, i = 0, 1, \dots, n + 1$ , where  $h = (b - a)/(n + 1)$

- We already have  $y_0 = u(a) = \alpha$  and  $y_{n+1} = u(b) = \beta$  from BC, and we seek approximate solution value  $y_i \approx u(t_i)$  at each interior mesh point  $t_i, i = 1, \dots, n$

Michael T. Heath

Scientific Computing

20 / 45

## Finite Difference Method, continued

- For these particular finite difference formulas, system to be solved is tridiagonal, which saves on both work and storage compared to general system of equations
- This is generally true of finite difference methods: they yield sparse systems because each equation involves few variables

Michael T. Heath

Scientific Computing

22 / 45

## Example, continued

- Replacing derivatives by standard finite difference approximations at  $t_1$  gives equation

$$\frac{y_2 - 2y_1 + y_0}{h^2} = f\left(t_1, y_1, \frac{y_2 - y_0}{2h}\right)$$

- Substituting boundary data, mesh size, and right hand side for this example we obtain

$$\frac{1 - 2y_1 + 0}{(0.5)^2} = 6t_1$$

or

$$4 - 8y_1 = 6(0.5) = 3$$

so that

$$y(0.5) \approx y_1 = 1/8 = 0.125$$

Michael T. Heath

Scientific Computing

24 / 45

## Example, continued

- In a practical problem, much smaller step size and many more mesh points would be required to achieve acceptable accuracy
- We would therefore obtain *system* of equations to solve for approximate solution values at mesh points, rather than single equation as in this example

< interactive example >



Michael T. Heath

Scientific Computing

25 / 45

## Collocation Method

- Popular choices of basis functions include polynomials, B-splines, and trigonometric functions
- Basis functions with global support, such as polynomials or trigonometric functions, yield *spectral method*
- Basis functions with highly localized support, such as B-splines, yield *finite element method*



Michael T. Heath

Scientific Computing

27 / 45

## Example: Collocation Method

- Consider again two-point BVP

$$u'' = 6t, \quad 0 < t < 1,$$

with BC

$$u(0) = 0, \quad u(1) = 1$$

- To keep computation to minimum, we use one interior collocation point,  $t = 0.5$
- Including boundary points, we have three collocation points,  $t_0 = 0$ ,  $t_1 = 0.5$ , and  $t_2 = 1$ , so we will be able to determine three parameters
- As basis functions we use first three monomials, so approximate solution has form

$$v(t, \mathbf{x}) = x_1 + x_2 t + x_3 t^2$$



Michael T. Heath

Scientific Computing

29 / 45

## Example, continued

- Solving this system of three equations in three unknowns gives

$$x_1 = 0, \quad x_2 = -0.5, \quad x_3 = 1.5$$

so approximate solution function is quadratic polynomial

$$u(t) \approx v(t, \mathbf{x}) = -0.5t + 1.5t^2$$

- At interior collocation point,  $t_2 = 0.5$ , we have approximate solution value

$$u(0.5) \approx v(0.5, \mathbf{x}) = 0.125$$



Michael T. Heath

Scientific Computing

31 / 45

## Collocation Method

- Collocation method* approximates solution to BVP by finite linear combination of basis functions
- For two-point BVP

$$u'' = f(t, u, u'), \quad a < t < b$$

with BC

$$u(a) = \alpha, \quad u(b) = \beta$$

we seek approximate solution of form

$$u(t) \approx v(t, \mathbf{x}) = \sum_{i=1}^n x_i \phi_i(t)$$

where  $\phi_i$  are basis functions defined on  $[a, b]$  and  $\mathbf{x}$  is  $n$ -vector of parameters to be determined



Michael T. Heath

Scientific Computing

26 / 45

## Collocation Method, continued

- To determine vector of parameters  $\mathbf{x}$ , define set of  $n$  *collocation points*,  $a = t_1 < \dots < t_n = b$ , at which approximate solution  $v(t, \mathbf{x})$  is forced to satisfy ODE and boundary conditions
- Common choices of collocation points include equally-spaced points or Chebyshev points
- Suitably smooth basis functions can be differentiated analytically, so that approximate solution and its derivatives can be substituted into ODE and BC to obtain system of algebraic equations for unknown parameters  $\mathbf{x}$



Michael T. Heath

Scientific Computing

28 / 45

## Example, continued

- Derivatives of approximate solution function with respect to  $t$  are given by

$$v'(t, \mathbf{x}) = x_2 + 2x_3 t, \quad v''(t, \mathbf{x}) = 2x_3$$

- Requiring ODE to be satisfied at interior collocation point  $t_2 = 0.5$  gives equation

$$v''(t_2, \mathbf{x}) = f(t_2, v(t_2, \mathbf{x}), v'(t_2, \mathbf{x}))$$

or

$$2x_3 = 6t_2 = 6(0.5) = 3$$

- Boundary condition at  $t_1 = 0$  gives equation

$$x_1 + x_2 t_1 + x_3 t_1^2 = x_1 = 0$$

- Boundary condition at  $t_3 = 1$  gives equation

$$x_1 + x_2 t_3 + x_3 t_3^2 = x_1 + x_2 + x_3 = 1$$




Michael T. Heath

Scientific Computing

30 / 45

## Example, continued



< interactive example >



Michael T. Heath

Scientific Computing

32 / 45

## Galerkin Method

- Rather than forcing residual to be zero at finite number of points, as in collocation, we could instead minimize residual over entire interval of integration
- For example, for *Poisson equation* in one dimension,

$$u'' = f(t), \quad a < t < b,$$

with homogeneous BC  $u(a) = 0, \quad u(b) = 0,$

substitute approx solution  $u(t) \approx v(t, \mathbf{x}) = \sum_{i=1}^n x_i \phi_i(t)$   
into ODE and define residual

$$r(t, \mathbf{x}) = v''(t, \mathbf{x}) - f(t) = \sum_{i=1}^n x_i \phi_i''(t) - f(t)$$

I

Michael T. Heath

Scientific Computing

33 / 45

## Galerkin Method, continued

- More generally, *weighted residual method* forces residual to be orthogonal to each of set of *weight functions* or *test functions*  $w_i$ ,

$$\int_a^b r(t, \mathbf{x}) w_i(t) dt = 0, \quad i = 1, \dots, n$$

- This yields linear system  $\mathbf{Ax} = \mathbf{b}$ , where now

$$a_{ij} = \int_a^b \phi_j''(t) w_i(t) dt, \quad b_i = \int_a^b f(t) w_i(t) dt$$

whose solution gives vector of parameters  $\mathbf{x}$

I

Michael T. Heath

Scientific Computing

35 / 45

## Galerkin Method, continued

- Degree of differentiability required can be reduced using integration by parts, which gives

$$\begin{aligned} \int_a^b v''(t, \mathbf{x}) \phi_i(t) dt &= v'(t) \phi_i(t) \Big|_a^b - \int_a^b v'(t) \phi_i'(t) dt \\ &= v'(b) \phi_i(b) - v'(a) \phi_i(a) - \int_a^b v'(t) \phi_i'(t) dt \end{aligned}$$

- Assuming basis functions  $\phi_i$  satisfy homogeneous boundary conditions, so  $\phi_i(0) = \phi_i(1) = 0$ , orthogonality condition then becomes

$$-\int_a^b v'(t) \phi_i'(t) dt = \int_a^b f(t) \phi_i(t) dt, \quad i = 1, \dots, n$$

I

Michael T. Heath

Scientific Computing

37 / 45


## Example: Galerkin Method

- Consider again two-point BVP

$$u'' = 6t, \quad 0 < t < 1,$$

with BC  $u(0) = 0, \quad u(1) = 1$

- We will approximate solution by piecewise linear polynomial, for which B-splines of degree 1 ("hat" functions) form suitable set of basis functions



- To keep computation to minimum, we again use same three mesh points, but now they become knots in piecewise linear polynomial approximation

I

Michael T. Heath

Scientific Computing

39 / 45

## Galerkin Method, continued

- Using *least squares method*, we can minimize

$$F(\mathbf{x}) = \frac{1}{2} \int_a^b r(t, \mathbf{x})^2 dt$$

by setting each component of its gradient to zero

- This yields symmetric system of linear algebraic equations  $\mathbf{Ax} = \mathbf{b}$ , where

$$a_{ij} = \int_a^b \phi_j''(t) \phi_i''(t) dt, \quad b_i = \int_a^b f(t) \phi_i''(t) dt$$

whose solution gives vector of parameters  $\mathbf{x}$

I

Michael T. Heath

Scientific Computing

34 / 45

## Galerkin Method, continued

- Matrix resulting from weighted residual method is generally not symmetric, and its entries involve second derivatives of basis functions

- Both drawbacks are overcome by *Galerkin method*, in which weight functions are chosen to be *same* as basis functions, i.e.,  $w_i = \phi_i, i = 1, \dots, n$

- Orthogonality condition then becomes

$$\int_a^b r(t, \mathbf{x}) \phi_i(t) dt = 0, \quad i = 1, \dots, n$$

or

$$\int_a^b v''(t, \mathbf{x}) \phi_i(t) dt = \int_a^b f(t) \phi_i(t) dt, \quad i = 1, \dots, n$$

I

Michael T. Heath

Scientific Computing

36 / 45

## Galerkin Method, continued

- This yields system of linear equations  $\mathbf{Ax} = \mathbf{b}$ , with

$$a_{ij} = -\int_a^b \phi_j'(t) \phi_i'(t) dt, \quad b_i = \int_a^b f(t) \phi_i(t) dt$$

whose solution gives vector of parameters  $\mathbf{x}$

- $\mathbf{A}$  is symmetric and involves only first derivatives of basis functions

I

Michael T. Heath

Scientific Computing

38 / 45

## Example, continued

- Thus, we seek approximate solution of form

$$u(t) \approx v(t, \mathbf{x}) = x_1 \phi_1(t) + x_2 \phi_2(t) + x_3 \phi_3(t)$$

- From BC, we must have  $x_1 = 0$  and  $x_3 = 1$

- To determine remaining parameter  $x_2$ , we impose Galerkin orthogonality condition on interior basis function  $\phi_2$  and obtain equation

$$-\sum_{j=1}^3 \left( \int_0^1 \phi_j'(t) \phi_2'(t) dt \right) x_j = \int_0^1 6t \phi_2(t) dt$$

or, upon evaluating these simple integrals analytically

$$2x_1 - 4x_2 + 2x_3 = 3/2$$

I

Michael T. Heath


Scientific Computing

40 / 45

## Example, continued

- Substituting known values for  $x_1$  and  $x_3$  then gives  $x_2 = 1/8$  for remaining unknown parameter, so piecewise linear approximate solution is

$$u(t) \approx v(t, \mathbf{x}) = 0.125\phi_2(t) + \phi_3(t)$$



- We note that  $v(0.5, \mathbf{x}) = 0.125$



## Eigenvalue Problems

- Standard eigenvalue problem for second-order ODE has form

$$u'' = \lambda f(t, u, u'), \quad a < t < b$$

with BC

$$u(a) = \alpha, \quad u(b) = \beta$$

where we seek not only solution  $u$  but also parameter  $\lambda$

- Scalar  $\lambda$  (possibly complex) is *eigenvalue* and solution  $u$  is corresponding *eigenfunction* for this two-point BVP
- Discretization of eigenvalue problem for ODE results in algebraic eigenvalue problem whose solution approximates that of original problem



## Example, continued

- Assuming  $g_i \neq 0$ , divide equation  $i$  by  $g_i$  for  $i = 1, \dots, n$ , to obtain linear system

$$A\mathbf{y} = \lambda\mathbf{y}$$

where  $n \times n$  matrix  $A$  has tridiagonal form

$$A = \frac{1}{h^2} \begin{bmatrix} -2/g_1 & 1/g_1 & 0 & \cdots & 0 \\ 1/g_2 & -2/g_2 & 1/g_2 & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & 1/g_{n-1} & -2/g_{n-1} & 1/g_{n-1} \\ 0 & \cdots & 0 & 1/g_n & -2/g_n \end{bmatrix}$$

- This standard algebraic eigenvalue problem can be solved by methods discussed previously



## Example, continued

- More realistic problem would have many more interior mesh points and basis functions, and correspondingly many parameters to be determined
- Resulting system of equations would be much larger but still sparse, and therefore relatively easy to solve, provided local basis functions, such as "hat" functions, are used
- Resulting approximate solution function is less smooth than true solution, but nevertheless becomes more accurate as more mesh points are used

< interactive example >



## Example: Eigenvalue Problem

- Consider linear two-point BVP

$$u'' = \lambda g(t)u, \quad a < t < b$$

with BC

$$u(a) = 0, \quad u(b) = 0$$

- Introduce discrete mesh points  $t_i$  in interval  $[a, b]$ , with mesh spacing  $h$  and use standard finite difference approximation for second derivative to obtain algebraic system

$$\frac{y_{i+1} - 2y_i + y_{i-1}}{h^2} = \lambda g_i y_i, \quad i = 1, \dots, n$$

where  $y_i = u(t_i)$  and  $g_i = g(t_i)$ , and from BC  $y_0 = u(a) = 0$  and  $y_{n+1} = u(b) = 0$

